



**NOAA  
FISHERIES**

**NW Fisheries  
Science Center**

# Overview of West Coast Stock Assessment Process and Staff

**Dr. Owen Hamel**  
**NWFSC Assessment Team Lead**

Disclaimer: This information is distributed solely for the purpose of pre-dissemination peer review under applicable information quality guidelines. It has not been formally disseminated by NOAA Fisheries. It does not represent and should not be construed to represent any agency determination or policy.

# Responsible for 90+ stocks in the Groundfish FMP

- This leads to some issues for stock assessment
  - Too many stocks for which to conduct complex assessments
    - given staff and review resource limitations
  - Insufficient data to conduct complex assessments for many stocks
  - Average catch approaches for setting catch limits not ideal




# History of Assessment Models used on U.S. West Coast

- Early variety of approaches
  - Stock Reduction Analysis (SRA)
  - Virtual Population Analysis (VPA) (cohort analysis)
  - Yield per Recruit Analysis (YPA)
- Later dominated by Statistical Catch at Age (CAA)
  - SS (SS1)
  - Independent models implemented in ADMB
  - SS2 – SS3 (in ADMB)



# How did we get here?: Evolution of fisheries models



Evolving towards more sophisticated, complex, & data-rich/needy models







# Advantages of Stock Synthesis

- Statistical catch-at-age models incorporate more realistic population dynamics
- Why Stock Synthesis?
  - Complex model which allows for a large variety of data types and model choices
  - New features can be added (as time allows)
  - Each version extensively tested and debugged
  - Shared R code for outputs (R4SS) and inputs
  - West coast assessment scientists have long history of working with SS and Rick Methot



# Scale, Status and Productivity



- Scale: Absolute level of biomass
- Status: Relative level of biomass
- Productivity: Natural rate of biomass change

# Population model

# Observation model

## Additional information


- Natural mortality
- Maturity
- Fecundity
- Spawner-recruit

## Fishery observations


- Landings and discards
- Biological samples

## Survey observations

- Catch
- Biological samples



# Progression of benchmark stock assessment models for the U.S. Pacific West Coast through 2011/2012






# Assessment Levels/Categories

## National Assessment Levels (SAIP 2011)

0. None
1. Index only
2. Simple life history equilibrium models
3. Aggregated production models
4. Size/age/stage-structured models
5. Space/Seasons/Ecosystem included

 = "Adequate"

## PFMC Assessment Categories


3. *Data-poor*
2. *Data-moderate* (includes some *benchmark* assessments)
1. *Data-rich* (includes most *benchmark* assessments)

# Ability to Assess 90+ stocks in the Groundfish FMP:

- Issues
  - Insufficient data to conduct complex assessments for many stocks
    - Trawl survey does not cover untrawlable habitat or nearshore areas
  - Insufficient resources to conduct complex assessments for others
  - Average catch approaches to setting catch limits not ideal
- Approach
  - Develop new methods
    - Data-poor – catch data + some life history
      - Simple, standard models; minimal review needed
    - Data-moderate – add in index of abundance
      - Intermediate complexity and level of review




# How did we get here?: (d)evolution of fisheries models



(D)evolving to models which require less data – but retain some of the features of more complex models

# How did we get here?: (d)evolution of fisheries models



(D)evolving to models which require less data – but retain some of the features of more complex models

- **And can be easily expanded with more data**

# Four Levels of Assessments

1. Benchmark (Full)
2. Update of benchmark
3. Data-Moderate
4. Data-Poor

# Four Levels of Assessments

1. Benchmark (Full)
2. Update of benchmark
3. Data-Moderate
4. Data-Poor

+ 3 more “assessment products”:

- Rebuilding Analyses
- Catch Reports
- Additional analyses requested by the PFMCM



# Benchmark Assessments

See C.2. for more information

- Typically use Stock Synthesis:
  - Allows for estimation of selectivity, natural mortality, productivity, recruitment, stock size; use of Bayesian priors, etc.
- Considerable exploration of uncertainty and model sensitivity
- Estimates of OFL (using MSY proxy) and status (% of SB0);
- Independent, interactive 4-5 day peer review of two assessments (STAR process), with final SSC review
- Used for 34 species since 2000



# Update Assessments

- Updates of prior Benchmark assessments
- New/revised data in previously used series; no new series
- Model structure remains the same
  - Previously estimated parameters remain estimated
  - Previously fixed parameters remain fixed
- Reviewed by SSC
  - Following initial review by SSC Groundfish Sub-committee





# Update Assessments

- Relatively simple to implement in terms of model and data choices;
  - i.e. essentially none
  - Previously fixed parameters remain fixed, previously estimated parameters remain estimated, etc.
  - No new data sources/types
- Problems can arise that cannot be dealt with within an update:
  - Poor fits to new or revised data
  - Odd parameter estimates
  - New and exciting data streams/types cannot be used
  - While new information on parameters/historical catch, etc. is allowed, no re-evaluation of other parameters.



# Data-Moderate Assessments

See C.4. for more information

- Recently-developed, intermediate between benchmark and data-poor
- Rely on catch data, 1 or more abundance indices, and assumptions about/priors on important parameters
- Methods underwent independent peer review (2012)
- 2013 assessments used STAR process;
  - Eventually will be reviewed by SSC Groundfish subcommittee
- Provide estimates of OFL and stock status
- “Adequate” assessments, developed/reviewed with fewer resources
- Successful initial use for 8 species in 2013



# Data-Poor Assessments

See C.3. for more information

- Simpler models with catch as only data input.
- Rely on prior distribution assumptions about important parameters:
  - current depletion ( $B_{\text{current}}/B_{\text{zero}}$ );
  - natural mortality rate ( $M$ );
  - $F_{\text{MSY}}/M$ ;
  - $B_{\text{MSY}}/B_{\text{zero}}$
- Produce uncertainty estimates, but model sensitivity not explored
- Methods underwent independent peer review (2011)
- Assessment results reviewed by SSC
- Provide information to set OFLs but not stock status;
- Generally a one-time analysis;
- Have been used for 50+ species since 2009.



# Assessment Adequacy

Relating SAIP to Council Criteria

NMFS Stock Assessment Improvement Plan	
Level Features	
1	CPUE only
2	Simple equilibrium models with life history; Yield-per-recruit; Catch curve
3	Production models w/ catch and index
4	Age-/length-based models
5	Add ecosystem elements or spatial/temporal parameter variation

PFMC Assessment Categories		
Tier	Features	Models used
3-low	Catch-only; no life history	
3-med	Aggregate catch and M	<b>Data Poor:</b> DCAC
3-high	Annual catch with M and age at 50% maturity	<b>Data Poor:</b> DB-SRA / SSS
2-low	M * survey biomass estimate	Rogers, '96
2-med	Historical catch and trend info	<b>Data Moderate:</b> XDB-SRA / exSSS
2-high	Age-/length-structured, but high uncertainty	<b>Full (STAR):</b> SS
1	Reliable age/length data; estimate year-class strength	<b>Full (STAR):</b> SS



# PFMC Assessment Categories

- SSC/Council assign assessments to Categories (1-3)
- Most **benchmark/update** assessments in **Category 1**
  - Those considered more uncertain may be placed in Category 2
- **Data-moderate** assessments in **Category 2**
- **Data-poor** assessments in **Category 3**
- Default scientific-uncertainty reductions from OFL to ABC
  - **Category 3 > Category 2 > Category 1**
- Assessment uncertainty may supersede default, if larger



# New Harvest Framework under MSA

OFL = Overfishing Level

ABC = Allowable Biological Catch

ACL = Annual catch limit

ACT = Annual catch target

Scientific  
uncertainty

Mgmt.  
uncertainty

$$\text{OFL} \geq \text{ABC} \geq \text{ACL} \geq \text{ACT}$$

$$\text{OFL} = F_{\text{MSY}} * B_{\text{Exploitable}}$$

$$\text{ABC} = F_{\text{MSY}} * B_{\text{EX}} * \text{Buffer}$$

(size of default buffer partially  
based on assessment category)



# Rebuilding Analyses

- Generally use stand-alone forward projection analysis program
  - SS configured to provide inputs
  - Prognosticates distribution of possible stock size under alternative harvesting schemes incorporating assessment model output on:
    - age composition at declaration and end year of assessment,
    - $M$ ,
    - fishery selectivity,
    - stock-recruit relationship
    - etc.
  - Provides catch projections, SPR, year of rebuild, uncertainty.
- XDB-SRA model has own rebuilding analysis approach



# Catch Reports

- Reporting of catch over recent years with best available data to ensure that catch levels are at or below the ACL.
  - An alternative to updates for rebuilding species with long rebuilding times and marginal indices of abundance.





# Additional Analyses for PFMC

- Groundfish Management Team often requests projections under alternative ACL/ACT or SPR values.
- Council Staff request projections from time to time as well.
  - 2013/2014 analysis for EIS:10-year forward projections across multiple harvest specifications and states of nature (base and alternative model parameterizations) for all stocks with assessments (even if a decade old).




# Current Staffing

- NWFSC
  - 9.7 Groundfish Stock Assessment Scientists
    - Tenure: a few days to ~15 years
      - Only one >7.5 years
  - 3 support staff
- SWFSC
  - 3 Groundfish Stock Assessment Scientists
  - 2 support staff



# History of Staffing Levels

# of Assessment Staff



# Staffing and Stock Assessments

## (Example: 2013)

- 11.7 Stock Assessment Scientists (Including 2 Team Leads) + 5 support staff
  - 8 Benchmark stock assessments in SS (Including Pacific hake)
    - 7 different lead authors
    - 1-2+ additional authors per assessment
  - 1 update assessment (in SS)
  - 9 Data-Moderate assessments (some with multiple areas)
    - 2 sets of 2 authors for the 8 reviewed in Data-Moderate STAR panel
    - 1 reviewed in STAR panel outside of Data-Moderate panel
  - 3 Catch Reports for rebuilding stocks
  - A few data-poor assessments performed late in the process
  - 1 rebuilding analysis
  - >> 400 10-year projections of 37 assessments for PFMC EIS analysis



# Evolution of Documentation

See B.2. for more information

- Simpler assessments documented methods and results along with needs for improvement in analysis.
- More complex assessments dependent on more diverse data led to increased need for documentation and review
- Terms of Reference with explicit document requirements
  - Executive summary with data used, assumptions/simplifications, main results and decision table.
  - Main document
  - Lists of tables and figures for display and diagnostics
  - Complete input files



# Evolution of Review Process

See D. for more information

- GMT review – prior to 1998
  - 3-4 Research Fishery Biologists and others
  - *More dependence upon subsequent SSC review*
- Stock Assessment Review (STAR) panels – 1998
  - Developed to provide thorough and independent peer review
  - Response to industry concerns about science
  - Full week panels with analyses being done each day/night
    - Issue with review process morphing into a workshop
      - Effort to refocus on being a review process while recognizing need to explore alternatives such that the results are best available science.



# Summary

- Stock assessment methods have evolved
- Complexity of benchmark assessments has increased in general
  - More common to have multiple authors than a decade ago
- Reductions in complexity of some assessments:
  - Use of data-moderate assessment methods
  - Simplifications in benchmark assessments when appropriate
- Assessment Category based upon:
  - assessment method
  - uncertainty in data and model outputs.
- NMFS staffing has increased a bit over time,
  - But so have quality of assessments, review and documentation,
  - Assessment contributions from state agencies/students have declined.



# Strengths

- Advanced stock assessment software, which allows for
  - Wide variety of data types
  - Multitude of modeling approaches
- Generalized R code for inputs and outputs
- Variety of assessment methods provides management advice for:
  - stocks with different amounts and types of data,
  - insufficient resources to conduct and review full assessment.
- Data from variety of surveys and fishery data collection programs
- Dedicated and innovative stock assessment scientists
- Comprehensive review process by highly qualified reviewers





# Challenges

- Data challenges:
  - Lack of coast-wide comprehensive survey in untrawlable habitat.
  - Inability to get accurate ages for a number of species
  - Surveys and fisheries generally stop at international boundaries
    - Fish do not
  - Delay in observing the size of a new recruitment
- Large number of species and small number of assessment scientists.
- Lack of stock assessment scientists from state agencies contributing
- Many demands on stock assessors time
- Many research projects to be conducted to improve stock assessment



# Solutions

- Data:
  - Develop visual, hook and line, pot or other surveys for untrawlable habitat
  - Continue to explore and improve ageing methods
  - Work to collaborate across borders
  - Continued improvement in data management, analysis and assessment.
  - Continue to explore environmental indices that correlate with recruitment
  - Look at other methods to estimate recruitment strength.
- Increase number of stock assessment scientists across both centers
- Collaborate within NMFS and NOAA, with academics and NGOs
- Encourage states to become more engaged in stock assessment.

